

Tumescent liposuction in lipoedema yields good long-term results

W. Schmeller, M. Hueppe* and I. Meier-Vollrath

Hanse-Klinik, St-Juergen-Ring 66, D-23564 Lübeck, Germany

*Department of Anaesthesiology, University of Lübeck, Ratzeburger Allee 160, D-23538 Lübeck, Germany

Summary

Correspondence

Wilfried Schmeller.

E-mail: ws@hanse-klinik.com

Accepted for publication

29 July 2011

Funding sources

None.

Conflicts of interest

None declared.

DOI 10.1111/j.1365-2133.2011.10566.x

Background Lipoedema is a painful disease in women with circumscribed increased subcutaneous fatty tissue, oedema, pain and bruising. Whereas conservative methods with combined decongestive therapy (manual lymphatic drainage, compression garments) have been well established over the past 50 years, surgical therapy with tumescent liposuction has only been used for about 10 years and long-term results are unknown.

Objectives To determine the efficacy of liposuction concerning appearance (body shape) and associated complaints after a long-term period.

Methods A total of 164 patients who had undergone conservative therapy over a period of years, were treated by liposuction under tumescent local anaesthesia with vibrating microcannulas. In a monocentric study, 112 could be re-evaluated with a standardized questionnaire after a mean of 3 years and 8 months (range 1 year and 1 month to 7 years and 4 months) following the initial surgery and a mean of 2 years and 11 months (8 months to 6 years and 10 months) following the last surgery.

Results All patients showed a distinct reduction of subcutaneous fatty tissue (average 9846 mL per person) with improvement of shape and normalization of body proportions. Additionally, they reported either a marked improvement or a complete disappearance of spontaneous pain, sensitivity to pressure, oedema, bruising, restriction of movement and cosmetic impairment, resulting in a tremendous increase in quality of life; all these complaints were reduced significantly ($P < 0.001$). Patients with lipoedema stage II and III showed better improvement compared with patients with stage I. Physical decongestive therapy could be either omitted (22.4% of cases) or continued to a much lower degree. No serious complications (wound infection rate 1.4%, bleeding rate 0.3%) were observed following surgery.

Conclusions Tumescent liposuction is a highly effective treatment for lipoedema with good morphological and functional long-term results.

Lipoedema, first described in the 1940s in the U.S.A.,^{1,2} is characterized by bilateral symmetric enlargement mainly of the legs as a result of abnormal deposition of subcutaneous fatty tissue in combination with oedema. Despite being a specified clinical entity, epidemiological data are still unknown.

The disease occurs exclusively in women; it is probably attributable to an autosomal dominant inheritance with sex limitation.³ In most cases, hips, thighs ('jodhpur-like riding breeches'), knees and lower legs, sometimes with a fatty cuff at the ankles (Turkish-pants phenomenon, inverse shouldering effect) are affected; arms are rarely affected and hands and feet

are never involved. The accumulation of fluid in the form of orthostatic oedema results in pain, tenderness and sensitivity to pressure; this is expressed in synonyms such as lipalgia, adiposalgia, adipopalgesia, adiposis dolorosa, lipomatosis dolorosa or painful column leg. Together with easy bruising, it causes significant physical morbidity.

Whereas lipoedema may appear in women with generalized obesity, body weight is normal in many patients. The obvious disproportion between a slim upper half of the body and large lower extremities cannot be eliminated by weight loss brought about by diet or physical exercise; this often results in considerable frustration and psychological problems.^{2,4}